

## Name changes and additions to the southern African freshwater fish fauna

PH Skelton

To cite this article: PH Skelton (2016): Name changes and additions to the southern African freshwater fish fauna, African Journal of Aquatic Science, DOI: [10.2989/16085914.2016.1186004](https://doi.org/10.2989/16085914.2016.1186004)

To link to this article: <http://dx.doi.org/10.2989/16085914.2016.1186004>



Published online: 16 Aug 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

## Name changes and additions to the southern African freshwater fish fauna

PH Skelton

South African Institute for Aquatic Biodiversity (SAIAB), Grahamstown, South Africa  
e-mail: [p.skelton@saiab.ac.za](mailto:p.skelton@saiab.ac.za)

**Changes made to the scientific names of southern African freshwater fishes since 2001 are explained and discussed. Adjustments to the phylogeny and classification of the fauna are outlined. Recent systematic studies on cyprinines are discussed and changes to the genera *Labeobarbus* (expanded concept), *Pseudobarbus* (expanded concept), and *Enteromius* are supported. The introduction of the family names Alestidae and Nothobranchiidae is discussed. Adjustments made to the genera *Nannocharax*, *Micropanchax*, *Tilapia*, *Coptodon* and *Mastacembelus* are explained. Species name changes for *Hepsetus* and *Zaireichthys* are detailed. New species described from the region, or resurrected from synonymy, since 2001 are listed.**

**Keywords:** classification, molecular DNA, nomenclature, phylogenetic relationships, phylogeny, systematics, taxonomy

The formal scientific names given to organisms are important for several reasons, firstly as a reliable and consistent means of communicating information about the organisms. Secondly, as expressions of scientific theory, the names are gateways to repositories of information about the organisms, their relationships and the natural order of life on earth. In 1758 Carolus Linnaeus bequeathed to humanity an enduring binomial system of formally naming plants and animals. In 1859, just one hundred years later, Charles Darwin provided an explanation and understanding of biological evolution that underpins the relationships and gives the logic for the natural order and classification of organisms, as embedded in their formal nomenclature. Taxonomic names are settled only once their specific status and interrelationships with other species are clearly understood. This knowledge frequently shifts as new species are discovered and new insights on relationships are revealed through research and the application of new techniques.

The advent of genetic and molecular DNA analysis has rejuvenated systematic biology in unprecedented ways over the past few decades. Digital technologies and approaches to the assembly and statistical analysis of large data sets have given further impetus to new knowledge and understandings of the interrelationships of organisms. As a consequence there has been a deluge of biodiversity discovery, sometimes in places where exploration was considered long complete or unproductive through the law of diminishing returns. The use of genetic data to identify organisms (DNA barcoding) has been introduced as a solution to deal with this unfolding discovery of new taxa. The need to establish as comprehensive a DNA reference library in the shortest possible time has itself driven the discovery of new biodiversity.

Fishes are relevant to humans in many ways and therefore are frequent subjects of scientific enquiry. In times

when names are changing frequently it is useful to provide an update and explanation for interested parties.

This note updates the changes to the names of southern African freshwater fishes since the publication of the keys and species accounts by Skelton (2001). It lists only species that have been formally described or names changed through scientific enquiry. It does not consider species known, but not yet formally described, of which there are many, as indicated in the IUCN list of species from southern Africa (Darwall et al. 2009).

In 2010 Wiley and Johnson (2010) published a teleost classification based on monophyletic groups. In the same year Chen and Mayden (2010) presented a 'phylogenomic perspective on the new era of ichthyology' that presaged a series of high-level studies using genomic analyses to determine the interrelationships of fishes. Near et al. (2012) provided the first multiple nuclear gene analysis across the spectrum of ray-finned fish diversity that adjusted the relationships for several groups where past resolution was uncertain. These include the sister-group relationships of the Ostariophysi with the Clupeomorph fishes, the relationships of salmoniform fishes and the reordering of several lineages leading to the 'superradiation' of spiny-rayed fishes. Near et al. (2013) provided a more detailed account of the 'superradiation' of spiny-rayed fishes or acanthomorphs. Two recent global level classifications of fishes based on these and other studies have been published on line under the 'DeepFin' Tree of Life banner (Betancur-R et al. 2013, 2014).

The classification by Wiley and Johnson (2010) made advances into disentangling the 'spiny-rayed bush' and extracted among others the mastacembelids and Cyprinodontiformes into a Series Smegmamorpharia. In the DeepFin classification (Betancur-R et al. 2014) these arrangements are adjusted again – so that the mastacembelids and

the anabantids are aligned within the series Anabantaria, whereas the Cyprinodontiformes are placed into the Series Ovalentaria, together with the cichlids, following Wainwright et al. (2012).

The name changes to genera and species have been recorded as follows:

1. Cyprinidae: Interrelationships within the large subfamily Cyprininae are complex, due in part to the fact that a high proportion of species (~400 species, >30%) are polyploid, either tetraploid, hexaploid or, in a few extreme cases, even octaploid. The discovery of polyploidy in African species of *Barbus* (Oellermann and Skelton 1990; Golubtsov and Krysanov 1993) attracted the attention of European researchers (Berrebi et al. 1996; Berrebi et al. 2014). Berrebi et al. (1996) proposed an initial division of the traditional genus *Barbus* along ploidy lines. The type species *Barbus barbatus* and related species in Europe are tetraploid, and the genus was restricted to this clade. The African hexaploid species were placed in *Labeobarbus*. However, the African diploid and tetraploid species were recognised as not belonging to this genus, and were designated as '*Barbus*' (Berrebi et al. 2014). The use of inverted commas here indicates uncertain placement, pending generic revision. Further resolution of the relationships of the African species came through molecular studies by Machordom and Doadrio (2001) and Tsigenopoulos et al. (2002, 2010).

The most detailed high-level molecular analysis of the cyprinine fishes to date (Yang et al. 2015), considered 117 genera and 612 species in four data sets, including five mitochondrial and cloned RAG1 nuclear genes. From the outcome the authors reclassified the Cyprininae into 11 tribes, 3 of them new, with relationships as indicated in Figure 1. African species are included in the tribes Labeonini, Torini and Smiliogastrini and the study confirmed that sub-Saharan African cyprinines are not closely related to European lineages. This therefore has nomenclatural implications for southern African species. Yang et al. (2015) suggested the following changes: (1) Hexaploid species previously included in *Barbus* be allocated to the genus *Labeobarbus*. The bounds of the genus *Labeobarbus* were expanded to include three large groups or lineages, the African labeobarbs, the '*Carasobarbus* lineage' from the Arabian peninsula and the '*Pterocapoeta* lineage' from the Levant and Middle East. (2) The tribe Smiliogastrini includes the minnows of Asia (*Puntius* and allies) as well as the diploid and tetraploid African species formerly included in *Barbus* or '*Barbus*'. The African tetraploid smiliogastrins are monophyletic, and so it was suggested that all these species be included in the genus *Pseudobarbus*, with the additional species tentatively as '*Pseudobarbus*'. The diploid African species formerly known as *Barbus* or '*Barbus*' were placed in the genus *Enteromius*.


The suggestions by Yang et al. (2015) have already been implemented in various studies (e.g. Skelton 2015; Armbruster et al. 2016; Decru et al. 2016; Vreven et al. 2016) and in the Catalogue of Fishes (Eschmeyer et al. n.d.). However, Schmidt and Bart (2015), in a letter to

the editor of 'Molecular Phylogenetics and Evolution', lodged objections and rejected the names suggested for African smiliogastrins. Consequently, certain authors (e.g. Ren and Mayden 2016; Stiassny et al. 2016; Stiassny and Sakharova 2016) have retained the use of "*Barbus*" or '*Barbus*' for African smiliogastrins. This taxonomic uncertainty is undesirable, and the decision to accept the suggestions by Yang et al. (2015) here is discussed further below.

The letter by Schmidt and Bart (2015) was titled 'Nomenclatural changes should not be based on equivocally supported phylogenies: Reply to Yang et al. 2015'. They accepted that Yang et al. provided 'reasonably well-supported hypotheses of relationships among 11 proposed tribes of cyprinine fishes' (Figure 1). However, they pointed out that support for the proposed relationships within some of the tribes was 'equivocal', i.e. they had less than 50% bootstrap support and that the results actually 'support multiple equally likely interpretations of relationships'. They pointed out that one clade within the smiliogastrins, including the genera *Systemus*, '*Barbus*' (*Enteromius*), *Barboides*, *Pseudobarbus* and *Clypeobarbus*, is well supported, but that 'collapsing branches with less than 50% support' resulted in a polytomy of the various intratribal clades. Therefore, they argued that the results within the smiliogastrine lineage were limited through taxon sampling, and that 'the poor resolution provides no support' for revalidation of the generic name *Enteromius* to accommodate all African diploid '*Barbus*'. They also rejected the suggestion designation by Yang et al. (2015) of '*Pseudobarbus*' for those southern African tetraploid barbs that don't fit into *Pseudobarbus* as it currently stands, and that a more apt treatment was to designate a new genus, or genera, for such species, or to revise *Pseudobarbus* appropriately.

African barbs are not closely related to *Barbus* (tribe Barbini) (Figure 1), and therefore the retention of the name *Barbus* for the African species is contrary to what a scientific genus name is intended to imply. To continue to use an interim, informal generic name linked to *Barbus*, i.e. '*Barbus*', does not solve the problem – it perpetuates it. Therefore, the African smiliogastrins require a more appropriate genus or genera within the tribe, and *Enteromius* is the earliest available name. Although the interrelationships of African smiliogastrins are relatively poorly understood, progress towards understanding these relationships is already in hand (Ren and Mayden 2016) and a more sustainable nomenclature is anticipated.

Yang et al. (2015) expand *Pseudobarbus* to include all tetraploid smiliogastrins from southern Africa, but indicated the generic uncertainties involved by placing the additional species in inverted commas as '*Pseudobarbus*'. The genus *Pseudobarbus* was defined by Skelton (1988) on morphological criteria, and included only the flexible dorsal rayed redfin minnows. Subsequently it was discovered that these species are tetraploid, as are certain other Cape cyprinids, including the cluster of 'serrated rayed' species from the region (Naran et al. 2006). The clustering by Yang et al. (2015) of southern African tetraploid cyprinids into



**Figure 1:** The relationships of cyprinine tribes, after Yang et al. (2015). African species are included in the tribes Labeonini, Torini and Smiliogastrini. The Barbini, including the genus *Barbus*, do not occur in Africa south of the Sahara

a well-supported clade is corroborated by molecular data from Tsigenopoulos et al. (2002, 2010), Berrebi et al. (2014) and unpublished molecular data (Swartz 2005; L da Costa, Lisbon, Portugal, pers. comm.). Within the framework of current African cyprinid taxonomy, an expanded genus *Pseudobarbus* for this clade, as suggested by Yang et al. (2015), is perfectly in order. Further deeper analysis of the lineage may well result in the newly added taxa being assigned to genera in their own right. Such is taxonomy!

Vreven et al. (2016) reviewed in detail the African tribe Torini, which includes the *Labeobarbus* and *Varicorhinus* species. Several significant taxonomic changes to southern African species emerged: (1) All the southern African *Varicorhinus* species were synonymised with *Labeobarbus*, which now includes *Labeobarbus dimidiatus*, *Labeobarbus nelspruitensis*, *Labeobarbus pungweensis* and *Labeobarbus xyrocheilus*. (2) The names of the Clanwilliam yellowfish (previously *Labeobarbus capensis*) and the Cape whitefish or witvis (previously *Barbus andrewi*) were changed to *Labeobarbus seeberi* (Gilchrist & Thompson, 1913), and '*Pseudobarbus*' *capensis* (Smith, 1841), respectively. These nomenclatural changes were made because the type specimen on which the name *Barbus*

*capensis* was based was re-identified as the Cape whitefish or witvis, and not the Clanwilliam yellowfish, as previously thought. The name '*Pseudobarbus*' *capensis* (Smith, 1841) for the Cape whitefish or witvis is in tune with the generic alignment of Yang et al. (2015). *Labeobarbus seeberi* (Gilchrist & Thompson, 1913) was resurrected as the earliest available name for the Clanwilliam yellowfish. In a detailed molecular phylogenetic study on the genus *Labeobarbus*, Beshera et al. (2016) confirm the relationships and realignment of *Varicorhinus nelspruitensis* to *Labeobarbus nelspruitensis*.

*Barbus bellcrossi* was included in the genus *Clypeobarbus* by Conway and Stiassny (2008). Stiassny and Sakharova (2016) provide a revised definition of the genus *Clypeobarbus*.

2. *Hepsetus*: Formerly, *Hepsetus odoe* was considered a monotypic genus distributed from the Senegal River to the upper Zambezi. The lineage has since been divided into six species, of which one, *Hepsetus cuvieri* Castelnau, 1861, occurs in southern Africa (Zengeya et al. 2011).
3. *Nannocharax*: The generic status of *Hemigrammocharax*, distinguished from *Nannocharax* on the dubious difference of an incomplete versus a complete lateral

**Table 1:** Additions to the list and changes to the names of southern African freshwater fishes since 2001. Species are presented in alphabetical order within systematic categories, according to the classification of Betancur-R (2014)

Former name	New name	New or resurrected species	Reference
<b>A. New species</b>			
		<i>Hippopotamyrus longilateralis</i>	Kramer and Swartz 2010
		<i>Hippopotamyrus szaboi</i>	Kramer et al. 2004
		<i>Marcusenius altisambesi</i>	Kramer et al. 2007
		<i>Marcusenius caudisquamatus</i>	Maake et al. 2014
		<i>Marcusenius desertus</i>	Kramer et al. 2016
		<i>Marcusenius krameri</i>	Maake et al. 2014
		<i>Marcusenius multisquamatus</i>	Kramer and Wink 2013
		<i>Marcusenius pongolensis</i>	Kramer et al. 2007
		<i>Petrocephalus longianalis</i>	Kramer et al. 2012
		<i>Petrocephalus longicapitis</i>	Kramer et al. 2012
		<i>Petrocephalus magnitrunci</i>	Kramer et al. 2012
		<i>Petrocephalus magnoculis</i>	Kramer et al. 2012
		<i>Petrocephalus okavangoensis</i>	Kramer et al. 2012
		<i>Petrocephalus petersi</i>	Kramer et al. 2012
		<i>Pollimyrus cuandoensis</i>	Kramer et al. 2014
		<i>Pollimyrus marianne</i>	Kramer et al. 2003
		<i>Pseudobarbus skeltoni</i>	Chakona and Swartz 2013
		<i>Pseudobarbus verlorenei</i>	Chakona et al. 2014
<i>Zaireichthys rotundiceps</i>		<i>Zaireichthys conspicuus</i>	Eccles et al. 2011
		<i>Zaireichthys kafuensis</i>	Eccles et al. 2011
<i>Zaireichthys rotundiceps</i>		<i>Zaireichthys kavangoensis</i>	Eccles et al. 2011
<i>Zaireichthys rotundiceps</i>		<i>Zaireichthys kunenensis</i>	Eccles et al. 2011
		<i>Zaireichthys monomotapia</i>	Eccles et al. 2011
		<i>Zaireichthys pallidus</i>	Eccles et al. 2011
		<i>Chetia brevicauda</i>	Bills and Weyl 2002
		<i>Nothobranchius boklundi</i>	Valdesalici 2010
		<i>Nothobranchius capriviensis</i>	Watters et al. 2015
<i>Nothobranchius rachovii</i> (in part)		<i>Nothobranchius kadleci</i>	Reichard 2010
<i>Nothobranchius rachovii</i> (in part)		<i>Nothobranchius krysanovi</i>	Schidlovskiy et al. 2010
		<i>Nothobranchius pienaari</i>	Schidlovskiy et al. 2010
<b>B. Resurrected species</b>			
<i>Marcusenius cubangoensis</i> Pellegrin, 1936		<i>Cyphomyrus cubangoensis</i>	Kramer and van der Bank 2011
<i>Fundulus mkuziensis</i> Fowler, 1934		<i>Nothobranchius mkuziensis</i>	Wildekamp 2004
<b>C. Renamed species</b>			
<i>Barbus bellcrossi</i>	<i>Clypeobarbus bellcrossi</i>		Conway and Stiassny 2008
<i>Barbus</i> spp.	<i>Enteromius</i> spp.		Yang et al. 2015
<i>Barbus andrewi</i>	' <i>Pseudobarbus</i> ' <i>capensis</i>		Yang et al. 2015; Vreven et al. 2016
<i>Barbus serra</i>	' <i>Pseudobarbus</i> ' <i>serra</i>		Yang et al. 2015
<i>Barbus calidus</i>	' <i>Pseudobarbus</i> ' <i>calidus</i>		Yang et al. 2015
<i>Barbus erubescens</i>	' <i>Pseudobarbus</i> ' <i>erubescens</i>		Yang et al. 2015
<i>Barbus hospes</i>	' <i>Pseudobarbus</i> ' <i>hospes</i>		Yang et al. 2015
<i>Barbus trevelyani</i>	' <i>Pseudobarbus</i> ' <i>trevelyani</i>		Yang et al. 2015
<i>Labeobarbus capensis</i>	<i>Labeobarbus seeberi</i>		Vreven et al. 2016
<i>Varicorhinus dimidiatus</i>	<i>Labeobarbus dimidiatus</i>		Vreven et al. 2016
<i>Varicorhinus nelspruitensis</i>	<i>Labeobarbus nelspruitensis</i>		Vreven et al. 2016
<i>Varicorhinus pungweensis</i>	<i>Labeobarbus pungweensis</i>		Vreven et al. 2016
<i>Varicorhinus xyrocheilus</i>	<i>Labeobarbus xyrocheilus</i>		Vreven et al. 2016
<i>Hemigrammocharax machadoi</i>	<i>Nannocharax machadoi</i>		Jerep and Vari 2013
<i>Hemigrammocharax multifasciatus</i>	<i>Nannocharax multifasciatus</i>		Jerep and Vari 2013
<i>Hepsetus odoe</i>	<i>Hepsetus cuvieri</i>		Zengeya et al. 2011
<i>Aethiomastacembelus</i> spp.	<i>Mastacembelus</i> spp.		Vreven 2005
<i>Tilapia rendalli</i>	<i>Coptodon rendalli</i>		Dunz and Schliwien 2013
<i>Aplocheilichthys hutereaui</i>	<i>Micropanchax hutereaui</i>		Ghedotti 2000
<i>Aplocheilichthys johnstoni</i>	<i>Micropanchax johnstoni</i>		Ghedotti 2000
<i>Aplocheilichthys katangae</i>	<i>Micropanchax katangae</i>		Ghedotti 2000
<i>Aplocheilichthys myaposae</i>	<i>Micropanchax myaposae</i>		Ghedotti 2000
<i>Aplocheilichthys</i> sp.	<i>Micropanchax</i> sp.		Ghedotti 2000

- line, was reviewed by Jerep and Vari (2013) and synonymised with *Nannocharax*. Two southern African species are affected by the outcome, viz. *Nannocharax machadoi* and *Nannocharax multifasciatus*.
4. Alestidae: The African characiform family Alestidae was regarded as monophyletic by Zanata and Vari (2005) and Calcagnotto et al. (2005) by using morphology and nuclear and mitochondrial gene sequences, respectively. Arroyave and Stiassny (2011) indicated that the family was not strictly monophyletic, because in their analysis two African genera, *Arnoldichthys* and *Lepidarchus*, were nested within Neotropical lineages. They suggested that the two genera be excluded from the family in order to maintain the monophyletic integrity of the Alestidae. These latter authors also found that, *inter alia*, the alestid genera *Brycinus* and *Rhabdalestes* were paraphyletic, but refrained from making taxonomic adjustments to the affected species, on the basis of these decisions being premature in the face of restricted taxon sampling.
  5. *Zaireichthys*: Skelton (2001) included two sand catlets of the genus *Zaireichthys* (*Z. rotundiceps* and *Z. sp.*) from southern Africa. Eccles et al. (2011) revised the genus and showed that *Z. rotundiceps* in southern Africa is a set of three species, *Zaireichthys kafuensis*, *Z. kunenensis* and *Z. monomotapa*. The second undescribed species in Skelton (2001) was described as *Z. pallidus* and two additional species new to the region were described as *Z. conspicuus* and *Z. kavangoensis*.
  6. Nothobranchiidae: Skelton (2001) used the family Aplocheilidae for the *Nothobranchius* killifishes. The classification of the family was updated by Lazara (2000), and consequently the appropriate family is now Nothobranchiidae Garman, 1895 (van der Laan et al. 2014).
  7. *Micropanchax*: The genus names of cyprinodontiform fishes are difficult to determine accurately, because the literature is a mixture of popular, semi-scientific and sound systematic scientific studies. The fishes are widespread, speciose and embrace challenging evolutionary, ecological and biological characteristics. Skelton (2001) used the genus *Aplocheilichthys* for the aplocheilichthyine topminnows. This genus is now restricted to a single species, *Aplocheilichthys spilauchen*, from coastal West Africa, and the other species are placed in the genus *Micropanchax* Myers, 1924, following the phylogenetic analysis and taxonomy of Ghedotti (2000).
  8. Cichlidae: Traditionally, the cichlids of East and southern Africa have been aligned in two clusters, the 'tilapiines' and the 'haplochromines', in accordance with morphological characters. Their manner of breeding (substrate spawning vs mouthbrooding) was important in defining the genera within each cluster. The mouthbrooding tilapiines were transferred to the genus *Oreochromis* by Trewavas (1983). Genetic studies have since provided new insight into African cichlid interrelationships, resetting the generic boundaries. Schwarzer et al. (2009) showed that the haplochromines and the tilapiines were completely integrated in a paraphyletic assemblage (conveniently called the haplotilapiines). Dunz and Schlieven (2013), assembling a large mitochondrial and nuclear gene data set from 76 species, untangled the haplotilapiine phylogeny sufficiently to reclassify the assemblage at the tribal and generic level. In southern Africa there are three tribes and genera: Coptodonini with the species *Coptodon rendalli*; Oreochromini with the genus *Oreochromis*; and Tilapiini with the genus *Tilapia*, which now contains only the southern African species *T. sparrmanii*, *T. baloni*, *T. ruweti* and *T. guinasana*.
  9. Mastacembelidae: Vreven (2005) found no justification for subdividing the African spiny-eels at generic level, and therefore the genus *Aethiomastacembelus*, as used by Skelton (2001), is discarded in favour of *Mastacembelus*.
- A total of 32 species have been described since 2001 (Table 1). Apart from the *Zaireichthys* sand catlets described by Eccles et al. (2011), as mentioned above, the new species are derived from only two families, 16 mormyrids, and six new species of *Nothobranchius* (family Nothobranchiidae). The mormyrid species were discovered by Kramer and his collaborators through studies on electric organ discharges, in combination with morphological and genetic characters.
- There is a sustained interest in *Nothobranchius* annual killifish, because of their extremophile life-history traits and attractiveness as aquarium species. One of the newly described species, *Nothobranchius capriviensis* Watters et al. (2015), refers to *Nothobranchius sp.* of Skelton (2001). *Nothobranchius rachovii* has been split into three different species (Schidlovskiy et al. 2010). Wildekamp (2004) resurrected *Nothobranchius mkuziensis* from, and placed *N. kuhntae* in synonymy with *N. othonotus*.
- Acknowledgements* — I appreciate and acknowledge the support given to me in my research by the Director, Dr Angus Paterson, and the staff and associates of SAIAB. The contributions of two anonymous reviewers considerably improved this article.

## References

- Armbruster JW, Stout CC, Hayes MM. 2016. An empirical test for convergence using African barbs (Cypriniformes: Cyprinidae). *Evolutionary Ecology* 30: 435–450.
- Arroyave J, Stiassny MLJ. 2011. Phylogenetic relationships and the temporal context for the diversification of African characins of the family Alestidae (Ostariophysi: Characiformes): evidence from DNA sequence data. *Molecular Phylogenetics and Evolution* 60: 385–397.
- Berrebi P, Chenuil A, Kotlik P, Machordom A, Tsigenopoulos CS. 2014. Disentangling the evolutionary history of the Genus *Barbus sensu lato*, a twenty years adventure. In: Alves MJ, Cartaxana A, Correia AM, Lopez LF (eds), *Professor Carlos Almaça (1934–2010) – Estado da Arte em Áreas Científicas que Desenvolveu*. Lisboa: Museu Nacional de História Natural e da Ciência. pp 29–55.
- Berrebi P, Kottelat M, Skelton P, Ráb P. 1996. Systematics of *Barbus*: state of the art and heuristic comments. *Folia Zoologica* 45: 5–12.
- Beshera KA, Harris PM, Mayden RL. 2016. Novel evolutionary lineages in *Labeobarbus* (Cypriniformes; Cyprinidae) based on phylogenetic analyses of mtDNA sequences. *Zootaxa* 4093: 363–381.
- Betancur-R R, Broughton RE, Wiley EO, Carpenter K, Lopez JA,

- Li C et al. 2013. The tree of life and a new classification of bony fishes. *PLoS Currents Tree of Life* 2013 April 18, Edition 1. Available at <http://currents.plos.org/treeoflife/article/the-tree-of-life-and-a-new-classification-of-bony-fishes/> [accessed 11 December 2015].
- Betancur-R R, Wiley EO, Bailly N, Miya M, Lecointre G, Orti G. 2014. Phylogenetic classification of bony fishes – Version 3. Available at [http://www.deepfin.org/Classification\\_v3.htm](http://www.deepfin.org/Classification_v3.htm) [accessed 11 December 2015].
- Bills IR, Weyl OLF. 2002. A new species of the Genus *Chetia* (Teleostei: Cichlidae) from the Lecitu River, Buzi System, Mozambique. *Smithiana Special Publication* 1: 1–11.
- Calcagnotto D, Schaefer SA, DeSalle R. 2005. Relationships among characiform fishes inferred from analysis of nuclear and mitochondrial gene sequences. *Molecular Phylogenetics and Evolution* 36: 135–153.
- Chakona A, Swartz ER. 2013. A new redfin species, *Pseudobarbus skeltoni* (Cyprinidae, Teleostei), from the Cape Floristic Region, South Africa. *Zootaxa* 3686: 565–577.
- Chakona A, Swartz ER, Skelton PH. 2014. A new species of redfin (Teleostei, Cyprinidae, *Pseudobarbus*), from the Verlorenvlei River system, South Africa. *ZooKeys* 453: 131–137.
- Chen W-J, Mayden RL. 2010. A phylogenomic perspective on the new era of ichthyology. *Bioscience* 60: 421–432.
- Conway KW, Stiassny MLJ. 2008. Phylogenetic diagnosis of the African cyprinid genus *Clypeobarbus* (Ostariophysi: Cyprinidae), with the rehabilitation of *Clypeobarbus bomokundi*. *Ichthyological Research* 55: 260–266.
- Darwall WRT, Smith KG, Tweddle D, Skelton PH (eds). 2009. *The status and distribution of freshwater biodiversity in southern Africa*. Gland: IUCN, and Grahamstown: SAIAB.
- Decru E, Moelants T, De Gelas K, Vreven E, Verheyen E, Snoeks J. 2016. Taxonomic challenges in freshwater fishes: a mismatch between morphology and DNA barcoding in fish of the north-eastern part of the Congo basin. *Molecular Ecology Resources* 16: 342–352.
- Dunz AR, Schliewen UK. 2013. Molecular phylogeny and revised classification of the haplotilapiine cichlid fishes formerly referred to as 'Tilapia'. *Molecular Phylogenetics and Evolution* 68: 64–80.
- Eschmeyer WN, Fricke R, van der Laan R (eds). n.d. Catalogue of fishes: genera, species, references. Available at <http://researcharchive.calacademy.org/research/ichthyology/catalogue/fishcatmain.asp> [accessed 3 March 2016]
- Eccles DH, Tweddle D, Skelton PH. 2011. Eight new species in the dwarf catfish genus *Zaireichthys* (Siluriformes: Amphiliidae). *Smithiana Bulletin* 13: 3–28.
- Ghedotti MJ. 2000. Phylogenetic analysis and taxonomy of the poecilioid fishes (Teleostei: Cyprinodontiformes). *Zoological Journal of the Linnean Society* 130: 1–53.
- Golubtsov AS, Krysanov EY. 1993. Karyological study of some cyprinid species from Ethiopia. The ploidy differences between large and small *Barbus* of Africa. *Journal of Fish Biology* 42: 445–455.
- Jerep FC, Vari RP. 2013. New species of *Hemigrammocharax* (Characiformes: Distichodontidae) from the northwestern Congo River Basin. *Copeia* 2013: 31–37.
- Kramer B, Bills IR, Skelton PH, Wink M. 2012. A critical revision of the churchill snoutfish, genus *Petrocephalus* Marcusen, 1854 (Actinopterygii: Teleostei: Mormyridae), from southern and eastern Africa, with the recognition of *Petrocephalus tanensis*, and the description of five new species. *Journal of Natural History* 46: 2179–2258.
- Kramer B, Skelton PH, van der Bank FH, Wink M. 2007. Allopatric differentiation in the *Marcusenius macrolepidotus* species complex in southern and eastern Africa: the resurrection of *M. pongolensis* and *M. angolensis*, and the description of two new species (Mormyridae, Teleostei). *Journal of Natural History* 41: 647–708.
- Kramer B, Swartz ER. 2010. A new species of slender stonebasher within the *Hippopotamyrus ansorgii* complex from the Cunene River in southern Africa (Teleostei: Mormyridae). *Journal of Natural History* 44: 2213–2242.
- Kramer B, van der Bank FH. 2011. The Victoria Falls, a species boundary for the Zambezi parrotfish, *Cyphomyrus discorhynchus* (Peters, 1852), and the resurrection of *Cyphomyrus cubangoensis* (Pellegrin, 1936) (Mormyridae: Teleostei). *Journal of Natural History* 45: 2669–2699.
- Kramer B, van der Bank FH, Flint N, Sauer-Gurth H, Wink M. 2003. Evidence for parapatric speciation in the mormyrid fish *Pollimyrus castelnaui* (Boulenger, 1911) from the Okavango-Upper Zambezi River systems: *P. marianne* sp. nov., defined by electric organ discharges, morphology and genetics. *Environmental Biology of Fishes* 67: 47–70.
- Kramer B, van der Bank FH, Wink M. 2004. The *Hippopotamyrus ansorgii* species complex in the Upper Zambezi River system with a description of a new species, *H. szaboi* (Mormyridae). *Zoologica Scripta* 33: 1–18.
- Kramer B, van der Bank H, Wink M. 2014. Marked differentiation in a new species of dwarf stonebasher, *Pollimyrus cuandoensis* sp. nov. (Mormyridae: Teleostei), from a contact zone with two sibling species of the Okavango and Zambezi rivers. *Journal of Natural History* 48: 429–463.
- Kramer LB, van der Bank FH, Wink M. 2016. *Marcusenius desertus* sp. nov. (Teleostei: Mormyridae), a mormyrid fish from the Namib desert. *African Journal of Aquatic Science* 2016: 1–18.
- Kramer B, Wink M. 2013. East-west differentiation in the *Marcusenius macrolepidotus* species complex in southern Africa: the description of a new species for the Cunene River, Namibia (Mormyridae, Teleostei). *Journal of Natural History* 47: 2327–2362.
- Lazara KJ. 2000. *The killifishes, an annotated checklist, synonymy and bibliography of recent oviparous cyprinodontiform fishes*. The Killifish Master Index 4. Cincinnati, Ohio: American Killifish Association, USA.
- Maake PA, Gon O, Swartz ER. 2014. Description of three new species of *Marcusenius* Gill, 1862 (Teleostei: Mormyridae) from South Africa and Mozambique. *Zootaxa* 3780: 455–480.
- Machordom A, Doadrio I. 2001. Evolutionary history and speciation modes in the cyprinid genus *Barbus*. *Proceedings of the Royal Society of London B* 268: 1297–1306.
- Naran D, Skelton PH, Villet MH. 2006. Karyology of the redfin minnows, genus *Pseudobarbus* Smith, 1841 (Teleostei: Cyprinidae): one of the evolutionary tetraploid lineages of South African barbines. *African Zoology* 41: 178–182.
- Near TJ, Eytan RI, Dornburg A, Kuhn KL, Moore JA, Davis MP et al. 2012. Resolution of ray-finned fish phylogeny and timing of diversification. *Proceedings of the National Academy of Sciences of the United States of America* 109: 13698–13703.
- Near TJ, Dornburg A, Eytan RI, Keck BP, Smith WL, Kuhn KL et al. 2013. Phylogeny and tempo of diversification in the superradiation of spiny-rayed fishes. *Proceedings of the National Academy of Sciences of the United States of America* 110: 12738–12743.
- Oellermann LK, Skelton PH. 1990. Hexaploidy in yellowfish species (*Barbus*, Pisces, Cyprinidae) from southern Africa. *Journal of Fish Biology* 37: 105–115.
- Reichard M. 2010. *Nothobranchius kadleci* (Cyprinodontiformes: Nothobranchiidae), a new species of annual killifish from central Mozambique. *Zootaxa* 2332: 49–60.
- Ren Q, Mayden RL. 2016. Molecular phylogeny and biogeography of African diploid barbs, '*Barbus*', and allies in Africa and Asia (Teleostei: Cypriniformes). *Zoologica Scripta*. DOI: 10.1111/zsc.12177 [accessed 25 March 2016].
- Schidlovskiy KM, Watters BR, Wildekamp RH. 2010. Notes on the annual killifish species *Nothobranchius rachovii*

- (Cyprinodontiformes; Nothobranchiidae) with the description of two new species. *Zootaxa* 2724: 37–57.
- Schmidt RC, Bart HL. 2015. Nomenclatural changes should not be based on equivocally supported phylogenies: Reply to Yang et al. 2015. *Molecular Phylogenetics and Evolution* 90: 193–194.
- Schwarzer J, Misof B, Tautz D, Schlieven UK. 2009. The root of the East African cichlid radiations. *BMC Evolutionary Biology* 3: 1–11.
- Skelton PH. 1988. A taxonomic revision of the redfin minnows (Pisces, Cyprinidae) from southern Africa. *Annals of the Cape Provincial Museums* 16: 201–307.
- Skelton PH. 2001. *A complete guide to the freshwater fishes of southern Africa*. Cape Town: Struik Publishers.
- Skelton PH. 2015. Fishes. In: Griffiths C, Day J, Picker M (eds), *Freshwater life: a field guide to the plants and animals of southern Africa*. Cape Town: Struik Nature. pp 96–123.
- Stiassny MLJ, Liyandja TLD, Monsembula Iyaba RJC. 2016. A new small barb (Cyprininae: Smiliogastrini) from the N'sele and Mayi Ndombe Rivers in the lower reaches of the Middle Congo Basin (Democratic Republic of Congo, Central Africa). *American Museum Novitates* 3848: 1–15.
- Stiassny MLJ, Sakharova H. 2016. Review of the smiliogastrin cyprinids of the Kwilu River (Kasai basin, Central Africa), revised diagnosis for *Clypeobarbus* (Cyprinidae: Cyprininae: Smiliogastrini) and description of a new species. *Journal of Fish Biology* 88: 1394–1412.
- Swartz ER. 2005. Phylogeography, phylogenetics and evolution of the redfins (Teleostei, Cyprinidae, *Pseudobarbus*) in southern Africa. PhD thesis, University of Pretoria, South Africa.
- Trewavas E. 1983. *Tilapiine fishes of the genera Sarotherodon, Oreochromis and Danakilia*. London: British Museum (Natural History).
- Tsigenopoulos CS, Ráb P, Naran D, Berrebi P. 2002. Multiple origins of polyploidy in the phylogeny of southern African barbs (Cyprinidae) as inferred from mtDNA markers. *Heredity* 88: 466–473.
- Tsigenopoulos CS, Kasapidis P, Berrebi P. 2010. Phylogenetic relationships of hexaploid large-sized barbs (genus *Labeobarbus*, Cyprinidae) based on mtDNA data. *Molecular Phylogenetics and Evolution* 56: 851–856.
- Valdesalici S. 2010. *Nothobranchius boklundi* (Cyprinodontiformes: Nothobranchiidae): a new annual killifish with two male colour morphs from the Luangwa River basin, Zambia. *Aqua, International Journal of Ichthyology* 16: 51–60.
- van der Laan R, Eschmeyer WN, Fricke R. 2014. Family-group names of recent fishes. *Zootaxa Monograph* 3882: 1–230. Available at <http://dx.doi.org/10.11646/zootaxa.3882.1.1> [accessed 3 March 2016].
- Vreven EJ. 2005. Mastacembelidae (Teleostei: Synbranchiformes) subfamily division and African generic division: an evaluation. *Journal of Natural History* 39: 351–370.
- Vreven EJWMN, Musschoot T, Snoeks J, Schlieven UK. 2016. The African hexaploid Torini (Cypriniformes: Cyprinidae): review of a tumultuous history. *Zoological Journal of the Linnean Society* 177: 231–305.
- Wainwright PC, Smith WL, Price SA, Tang KL, Sparks JS, Ferry LA et al. 2012. The evolution of pharyngognath: a phylogenetic and functional appraisal of the pharyngeal jaw key innovation in labroid fishes and beyond. *Systematic Biology* 61: 1001–1027.
- Watters BR, Wildekamp RH, Shidlovskiy KM. 2015. Description and biogeography of *Nothobranchius caprivensis*, a new species of annual killifish from the Zambezi Region of Namibia (Cyprinodontiformes: Nothobranchiidae). *Journal of the American Killifish Association* 47: 97–133.
- Wildekamp RH. 2004. *A world of killies. Atlas of the oviparous cyprinodontiform fishes of the world*. Vol. IV. Elyria, Ohio: American Killifish Association.
- Wiley EO, Johnson GD. 2010. A teleost classification based on monophyletic groups. In: Nelson JS, Schultz HP, Wilson MVH (eds), *Origin and phylogenetic interrelationships of teleosts*. Munich: Verlag Dr Friedrich Pfeil. pp 123–182.
- Yang L, Sado T, Vincent Hirt M, Pasco-Viel E, Arunachalam M, Li J et al. 2015. Phylogeny and polyploidy: resolving the classification of cyprinine fishes (Teleostei: Cypriniformes). *Molecular Phylogenetics and Evolution* 85: 97–116.
- Zanata AM, Vari RP. 2005. The family Alestidae (Ostariophysi, Characiformes): a phylogenetic analysis of a trans-Atlantic clade. *Zoological Journal of the Linnean Society* 145: 1–144.
- Zengeya, TA, Decru E, Vreven EJ. 2011. Revalidation of *Hepsetus cuvieri* (Castelnau, 1861) (Characiformes: Hepsetidae) from the Quanza, Zambezi and southern part of the Congo ichthyofauna provinces. *Journal of Natural History* 45: 1723–1744.